



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

EIF4G1 (Human) Recombinant Protein (Q01)

Alternative splicing results in five transcript variants encoding four distinct isoforms. [provided by RefSeq]

Catalog Number: H00001981-Q01

Regulation Status: For research use only (RUO)

Product Description: Human EIF4G1 partial ORF (NP_886553, 1500 a.a. - 1599 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

DVAVLKARAKLLQKYLQDEQKELQALYALQALVVTLEQ
PPNLLRMFFDALYDEDVVKEDAFYSWESSKDPAEQQ
GKGVALKSVTAFFKWLREAEESDHN

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 36.74

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1981

Gene Symbol: EIF4G1

Gene Alias: DKFZp686A1451, EIF4F, EIF4G, p220

Gene Summary: The protein encoded by this gene is a component of the protein complex EIF4F, which is involved in the recognition of the mRNA cap, ATP-dependent unwinding of 5'-terminal secondary structure, and recruitment of mRNA to the ribosome.