



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

F3 (Human) IP-WB Antibody Pair

Catalog # : H00002152-PW3

規格 : [1 Set]


List All

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of F3 transfected lysate using rabbit polyclonal anti-F3 and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with mouse purified polyclonal anti-F3.

Supplied Product: Antibody pair set content:
 1. Antibody pair for IP: rabbit polyclonal anti-F3 (300 ul)
 2. Antibody pair for WB: mouse purified polyclonal anti-F3 (50 ug)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [2152](#)

Gene Name: F3

Gene Alias: CD142,TF,TFA

Gene Description: coagulation factor III (thromboplastin, tissue factor)

Omim ID: [134390](#)


Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes coagulation factor III which is a cell surface

glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. [provided by RefSeq

Other Designations: OTTHUMP00000012426,coagulation factor III,tissue factor

Interactome



Gene Pathway

[Complement and coagulation cascades](#)

Related Disease

[Acute Disease](#) [Alzheimer Disease](#) [Alzheimer disease](#) [Atherosclerosis](#) [Atherosclerosis](#)
[Brain Ischemia](#) [Cardiovascular Diseases](#) [Carotid Stenosis](#) [Chorioamnionitis](#)
[Coronary Thrombosis](#) [Diabetes Complications](#) [Diabetes Mellitus, Type 2](#) [Edema](#)
[Fetal Membranes, Premature Rupture](#) [Genetic Predisposition to Disease](#) [Hemorrhage](#)
[Kidney Failure, Chronic](#) [Metabolic Syndrome X](#) [Myocardial Ischemia](#)

[... see more](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.