

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





F11 (Human) IP-WB Antibody Pair

Catalog #: H00002160-PW1 規格:[1 Set]

List All

Specification		Application Image
Product Description:	This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.	Immunoprecipitation-Western Blot
Reactivity:	Human	
Quality Control Testing:	Immunoprecipitation-Western Blot (IP-WB)	
	150 - 100 -	
	75 -	
	50 - 37 -	
	25-	
	20-	
	15=	
	Immunoprecipitation of F11 transfected lysate using mouse monoclonal anti-F11 and Protein A Magnetic Bead (<u>U0007</u>), and immunoblotted with rabbit polyclonal anti-F11.	
Supplied Product:	Antibody pair set content: 1. Antibody pair for IP: mouse monoclonal anti-F11 (300 ug) 2. Antibody pair for WB: rabbit polyclonal anti-F11 (50 ul)	
Storage Instruction:	Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.	
MSDS:	<u>Download</u>	
Applications		
Immunoprecipit	ation-Western Blot	
Gene Informatio	on	
Entrez GenelD:	2160	
Gene Name:	F11	
Gene Alias:	FXI,MGC141891	
Gene Description:	coagulation factor XI	
Omim ID:	264900	

Page 1 of 2 2016/5/20

Gene Summary: This gene encodes coagulation factor XI of the blood coagulation cascade. This protein is present in plasma as a zymogen, which is a unique plasma coagulation enzyme because it exists as a homodimer consisting of two identical polypeptide chains linked by disulfide bonds. During activation of the plasma factor XI, an internal peptide bond is cleaved by factor XIIa (or XII) in each of the two chains, resulting in activated factor XIa, a serine protease composed of two heavy and two light chains held together by disulfide bonds. This activated plasma factor XI triggers the middle phase of the intrisic pathway of blood coagulation by activating factor IX. Defects in this factor lead to Rosenthal syndrome, a blood coagulation abnormality. [provided by RefSeq

Other **Designations:** plasma thromboplastin antecedent, platelet coagulation factor XI

Gene Pathway

Complement and coagulation cascades

Related Disease

<u>Cardiovascular Diseases</u> <u>Diabetes Mellitus</u>, <u>Type 2 Edema Factor XI Deficiency</u> Genetic Predisposition to Disease Hypertension, Pregnancy-Induced Kidney Failure, Chronic Pre-Eclampsia Thrombophilia Venous Thrombosis

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 2 of 2 2016/5/20