



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FCAR 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00002204-T02

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-FCAR full-length

Host: Human

Theoretical MW (kDa): 32.3


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-FCAR antibody (H00002204-B02) by Western Blots.

SDS-PAGE Gel



FCAR transfected lysate.

Western Blot



Lane 1: FCAR transfected lysate (32.3 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [2204](#)

GeneBank Accession#: [NM_002000.2](#)

Protein Accession#: [NP_001991.1](#)

Gene Name: FCAR

Gene Alias: CD89

Gene Description: Fc fragment of IgA, receptor for

Omim ID: [147045](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsonized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq]

Other Designations: Fc alpha receptor, IgA Fc receptor, OTTHUMP00000068692, OTTHUMP00000068696, OTTHUMP00000068697, OTTHUMP00000068698

Related Disease

[Aggressive Periodontitis](#) [Alveolar Bone Loss](#) [Asthma](#) [Asthma](#) [Brain Ischemia](#) [Cardiovascular Diseases](#) [Chronic Periodontitis](#) [Coronary Disease](#) [Genetic Predisposition to Disease](#) [Glomerulonephritis](#) [IGA](#) [Hepatitis C](#) [Chronic Lichen Planus](#) [Oral Lupus Erythematosus](#) [Systemic Myocardial Infarction](#) [Periodontal Attachment Loss](#) [Periodontal Pocket](#) [Periodontitis](#) [Stroke](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.