



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## FGFR2 Pre-design Chimera RNAi

Catalog # : H00002263-R51

規格 : [ 10 nmol ] [ 20 nmol ]

List All

### Specification

**Product Description:** Homo sapiens fibroblast growth factor receptor 2 (bacteria-expressed kinase, keratinocyte growth factor receptor, craniofacial dysostosis 1, Crouzon syndrome, Pfeiffer syndrome, Jackson-Weiss syndrome) (FGFR2), transcript variant 12, mRNA.

**Reactivity:** Human


**Supplied Product:** DEPC water

**Target Refseq:** NM\_023030

**Target Region:** Coding sequence

**Storage Instruction:** Store at -20°C, do not exceed 4 - 5 freeze-thaw cycles to ensure product integrity.

**Note:** Position of the Chimera RNAi.  
The related RNAi products listed below were designed from different accession number but sharing the same RNAi sequence.



### Publication Reference

- [dsCheck: highly sensitive off-target search software for double-stranded RNA-mediated RNA interference.](#)  
Naito Y, Yamada T, Matsumiya T, Ui-Tei K, Saigo K, Morishita S. *Nucleic Acids Res.* 2005 Jul 1;33(Web Server issue):W589-91.
- [Functional dissection of siRNA sequence by systematic DNA substitution: modified siRNA with a DNA seed arm is a powerful tool for mammalian gene silencing with significantly reduced off-target effect.](#)  
Ui-Tei K, Naito Y, Zenno S, Nishi K, Yamato K, Takahashi F, Juni A, Saigo K. *Nucleic Acids Res.* 2008 Apr;36(7):2136-51. Epub 2008 Feb 11.
- [Guidelines for the selection of highly effective siRNA sequences for mammalian and chick RNA interference.](#)  
Ui-Tei K, Naito Y, Takahashi F, Haraguchi T, Ohki-Hamazaki H, Juni A, Ueda R, Saigo K. *Nucleic Acids Res.* 2004 Feb 9;32(3):936-48. Print 2004.
- [siDirect: highly effective, target-specific siRNA design software for mammalian RNA interference.](#)  
Naito Y, Yamada T, Ui-Tei K, Morishita S, Saigo K. *Nucleic Acids Res.* 2004 Jul 1;32(Web Server issue):W124-9.

### Applications

RNAi Knockdown

### Gene Information

Entrez GeneID: [2263](#)

### Application Image

RNAi Knockdown

**Gene Name:** FGFR2

**Gene Alias:** BEK,BFR-1,CD332,CEK3,CFD1,ECT1,FLJ98662,JWS,K-SAM,KGFR,TK14,TK25

**Gene Description:** fibroblast growth factor receptor 2


**Omim ID:** [101200](#), [101400](#), [101600](#), [123150](#), [123500](#), [123790](#), [137215](#), [149730](#), [176943](#), [207410](#)

**Gene Ontology:** [Hyperlink](#)

**Gene Summary:** The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member is a high-affinity receptor for acidic, basic and/or keratinocyte growth factor, depending on the isoform. Mutations in this gene are associated with Crouzon syndrome, Pfeiffer syndrome, Craniosynostosis, Apert syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrata syndrome, Saethre-Chotzen syndrome, and syndromic craniosynostosis. Multiple alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq]

**Other Designations:** BEK fibroblast growth factor receptor,FGF receptor,OTTHUMP00000020621,OTTHUMP00000020629,bacteria-expressed kinase,hydroxyaryl-protein kinase,keratinocyte growth factor receptor,protein tyrosine kinase, receptor like 14,soluble FGFR4 variant 4

## Interactome



## Gene Pathway

[Endocytosis](#) [MAPK signaling pathway](#) [Pathways in cancer](#) [Prostate cancer](#)  
[Regulation of actin cytoskeleton](#)

## Related Disease

[Acrocephalosyndactylia](#) [Adenocarcinoma](#) [Adenocarcinoma, Mucinous](#) [Alzheimer Disease](#)  
[Alzheimer disease](#) [Birth Weight](#) [Breast cancer](#) [Breast Neoplasms](#) [Bronchial Hyperreactivity](#)  
[Carcinoma, Basal Cell](#) [Carcinoma, Endometrioid](#) [Carcinoma, Squamous Cell](#)  
[Cardiovascular Diseases](#) [Cleft Lip](#) [Cleft Palate](#) [Craniofacial Dysostosis](#) [Craniosynostoses](#)  
[Cystadenocarcinoma, Serous](#) [Depressive Disorder, Major](#)

[... see more](#)

---

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.