



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FGR 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00002268-T01

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-FGR full-length

Host: Human

Theoretical MW (kDa): 58.3


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-FGR antibody (H00002268-B01) by Western Blots.

SDS-PAGE Gel



FGR transfected lysate.

Western Blot



Lane 1: FGR transfected lysate (58.3 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [2268](#)

GeneBank Accession#: [NM_005248.2](#)

Protein Accession#: [NP_005239.1](#)

Gene Name: FGR

Gene Alias: FLJ43153,MGC75096,SRC2,c-fgr,c-src2,p55c-fgr,p58c-fgr

Gene Description: Gardner-Rasheed feline sarcoma viral (v-fgr) oncogene homolog

Omim ID: [164940](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein contains N-terminal sites for myristylation and palmitoylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to plasma membrane ruffles, and functions as a negative regulator of cell migration and adhesion triggered by the beta-2 integrin signal transduction pathway. Infection with Epstein-Barr virus results in the overexpression of this gene. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq]

Other Designations: OTTHUMP00000003601,OTTHUMP00000003602,OTTHUMP00000003603,c-fgr protooncogene,c-src-2 proto-oncogene,p55-c-fgr protein,proto-oncogene tyrosine-protein kinase FGR

Gene Pathway

[Chemokine signaling pathway](#)

Related Disease

[HIV Infections](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.