



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## GTF2I 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00002969-T01

規格 : [ 100 uL ]

List All

### Specification

**Transfected Cell Line:** 293T


**Plasmid:** pCMV-GTF2I full-length

**Host:** Human

**Theoretical MW (kDa):** 107.47


**Quality Control Testing:** Transient overexpression cell lysate was tested with Anti-GTF2I antibody ([H00002969-B01](#)) by Western Blots.

#### SDS-PAGE Gel



GTF2I transfected lysate.

#### Western Blot



Lane 1: GTF2I transfected lysate ( 107.47 KDa)

Lane 2: Non-transfected lysate.

**Storage Buffer:** 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

**Storage Instruction:** Store at -80°C. Aliquot to avoid repeated freezing and thawing.

**MSDS:**  [Download](#)

### Applications

## Western Blot

### Gene Information

**Entrez GeneID:** [2969](#)

**GeneBank Accession#:** [BC099907.1](#)

**Protein Accession#:** [AAH99907.1](#)

**Gene Name:** GTF2I

**Gene Alias:** BAP-135,BAP135,BTKAP1,DIWS,FLJ38776,FLJ56355,IB291,SPIN,TFII-I,WBS,WBSCR6

**Gene Description:** general transcription factor II, i

**Omim ID:** [194050](#), [601679](#)

**Gene Ontology:** [Hyperlink](#)

**Gene Summary:** This gene encodes a multifunctional phosphoprotein with roles in transcription and signal transduction. It is deleted in Williams-Beuren syndrome, a multisystem developmental disorder caused by the deletion of contiguous genes at chromosome 7q11.23. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 7, 13 and 21. [provided by RefSeq]

**Other Designations:** BTK-associated protein, 135kD,Bruton tyrosine kinase-associated protein 135,Williams-Beuren syndrome chromosome region 6

### Gene Pathway

[Basal transcription factors](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.