



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HABP2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00003026-T01

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-HABP2 full-length

Host: Human

Theoretical MW (kDa): 61.71


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-HABP2 antibody ([H00003026-B01](#)) by Western Blots.

SDS-PAGE Gel



HABP2 transfected lysate.

Western Blot



Lane 1: HABP2 transfected lysate (61.71 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [3026](#)

GeneBank [NM_004132.2](#)
Accession#:

Protein [NP_004123.1](#)
Accession#:

Gene Name: HABP2

Gene Alias: FSAP,HABP,HGFAL,PHBP

Gene Description: hyaluronan binding protein 2

Omim ID: [188050](#), [603924](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is an extracellular serine protease that binds hyaluronic acid and is involved in cell adhesion. The encoded protein is synthesized as a single chain, but then undergoes an autoproteolytic event to form the functional heterodimer. Further autoproteolysis leads to smaller, inactive peptides. This protease is known to cleave urinary plasminogen activator, coagulation factor VII, and the alpha and beta chains of fibrinogen, but not prothrombin, plasminogen, or the gamma chain of fibrinogen. [provided by RefSeq]

Other Designations: OTTHUMP00000020505, factor VII activating protein, hepatocyte growth factor activator-like protein, hyaluronic acid binding protein 2, plasma hyaluronan binding protein

Related Disease

[Acute Disease](#) [Alzheimer Disease](#) [Arterial Occlusive Diseases](#) [Atherosclerosis](#)
[Cardiovascular Diseases](#) [Diabetes Mellitus, Type 2](#) [Disease Progression](#) [Edema](#)
[Genetic Predisposition to Disease](#) [Hemorrhagic Disorders](#) [Hepatitis C](#) [Liver Cirrhosis](#)
[Thromboembolism](#) [Thrombophilia](#) [Venous Thrombosis](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.