



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

HADH (Human) IP-WB Antibody Pair

Catalog # : H00003033-PW2

規格 : [1 Set]


[List All](#)

Specification

Product Description: This IP-WB antibody pair set comes with one antibody for immunoprecipitation and another to detect the precipitated protein in western blot.

Reactivity: Human

Quality Control Testing: Immunoprecipitation-Western Blot (IP-WB)



Immunoprecipitation of HADH transfected lysate using mouse monoclonal anti-HADH and Protein A Magnetic Bead ([U0007](#)), and immunoblotted with rabbit polyclonal anti-HADH.

Supplied Product: Antibody pair set content:
1. Antibody pair for IP: mouse monoclonal anti-HADH (300 ug)
2. Antibody pair for WB: rabbit polyclonal anti-HADH (50 ul)

Storage Instruction: Store reagents of the antibody pair set at -20°C or lower. Please aliquot to avoid repeated freeze thaw cycle. Reagents should be returned to -20°C storage immediately after use.

MSDS:  [Download](#)

Applications

Immunoprecipitation-Western Blot

 [Protocol Download](#)

Gene Information

Entrez GeneID: [3033](#)

Gene Name: HADH

Gene Alias: HAD, HADH1, HADHSC, HHF4, M/SCHAD, MGC8392, SCHAD

Gene Description: hydroxyacyl-Coenzyme A dehydrogenase

Omim ID: [231530](#), [601609](#), [609975](#)

Gene Ontology: [Hyperlink](#)

Application Image

Immunoprecipitation-Western Blot

Gene Summary: This gene is a member of the 3-hydroxyacyl-CoA dehydrogenase gene family. The encoded protein functions in the mitochondrial matrix to catalyze the oxidation of straight-chain 3-hydroxyacyl-CoAs as part of the beta-oxidation pathway. Its enzymatic activity is highest with medium-chain-length fatty acids. Mutations in this gene cause one form of familial hyperinsulinemic hypoglycemia. The human genome contains a related pseudogene. [provided by RefSeq]

Other Designations: L-3-hydroxyacyl-Coenzyme A dehydrogenase, L-3-hydroxyacyl-Coenzyme A dehydrogenase, short chain

Gene Pathway

[Butanoate metabolism](#) [Caprolactam degradation](#) [Fatty acid elongation in mitochondria](#) [Fatty acid metabolism](#) [Geraniol degradation](#) [Lysine degradation](#) [Metabolic pathways](#) [Tryptophan metabolism](#) [Valine, leucine and isoleucine degradation](#)

Related Disease

[Alcoholism](#) [Diabetes Mellitus, Type 2](#) [Genetic Predisposition to Disease](#) [Hyperinsulinism](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.