



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

NRG1 (Human) Recombinant Protein (P01)

Catalog Number: H00003084-P01

Regulation Status: For research use only (RUO)

Product Description: Human NRG1 full-length ORF (NP_039252.1, 1 a.a. - 241 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MSEKKEGRGKGGKGGKKKERGSGKKPESAAGSQSPALP
PQLKEMKSQESAAGSKLVLCETSSEYSSLRFKWFKN
GNELNRKNKPQNIKIQQKPGKSELRINKASLADSGEYM
CKVISKLGNDASANITIVESNEITGMPASTEGAYVSS
ESPIRISVSTEGANTSSSTSTSTTGTSHLVKCAEKEKTF
CVNGGECFMVKDLSNPSRYLCKCPNEFTGDRCQNYV
MASFYSTSTPFLSLPE

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 52.5

Applications: AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3084

Gene Symbol: NRG1

Gene Alias: ARIA, GGF, GGF2, HGL, HRG, HRG1, HRGA, NDF, SMDF

Gene Summary: The protein encoded by this gene was originally identified as a 44-kD glycoprotein that interacts with the NEU/ERBB2 receptor tyrosine kinase to increase its phosphorylation on tyrosine residues. This protein is a signaling protein that mediates cell-cell interactions and plays critical roles in the growth and development of multiple organ systems. It is known that an extraordinary variety of different isoforms are produced from this gene through alternative promoter usage and splicing. These isoforms are tissue-specifically expressed and differ significantly in their structure, and thereby these isoforms are classified into types I, II, III, IV, V and VI. The gene dysregulation has been linked to diseases such as cancer, schizophrenia and bipolar disorder (BPD). [provided by RefSeq]