



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## HK2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00003099-T01

規格 : [ 100 uL ]

[List All](#)

### Specification

**Transfected Cell Line:** 293T


**Plasmid:** pCMV-HK2 full-length

**Host:** Human

**Theoretical MW (kDa):** 100.98


**Quality Control Testing:** Transient overexpression cell lysate was tested with Anti-HK2 antibody (H00003099-B01) by Western Blots.

#### SDS-PAGE Gel



HK2 transfected lysate.

#### Western Blot



Lane 1: HK2 transfected lysate ( 100.98 KDa)

Lane 2: Non-transfected lysate.

**Storage Buffer:** 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

**Storage Instruction:** Store at -80°C. Aliquot to avoid repeated freezing and thawing.

**MSDS:**  [Download](#)

### Applications

## Western Blot

### Gene Information

Entrez GeneID: [3099](#)

GeneBank Accession#: [NM\\_000189](#)

Protein Accession#: [NP\\_000180](#)

Gene Name: HK2

Gene Alias: DKFZp686M1669, HKII, HXK2

Gene Description: hexokinase 2

Omim ID: [601125](#)

Gene Ontology: [Hyperlink](#)

**Gene Summary:** Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes hexokinase 2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-responsive, and studies in rat suggest that it is involved in the increased rate of glycolysis seen in rapidly growing cancer cells. [provided by RefSeq]

Other Designations: hexokinase-2, muscle

### Gene Pathway

[Amino sugar and nucleotide sugar metabolism](#)

[Biosynthesis of alkaloids derived from histidine and purine](#)

[Biosynthesis of alkaloids derived from ornithine, lysine and nicotinic acid](#)

[Biosynthesis of alkaloids derived from shikimate pathway](#)

[Biosynthesis of alkaloids derived from terpenoid and polyketide](#)

[Biosynthesis of phenylpropanoids](#) [Biosynthesis of plant hormones](#)

[Biosynthesis of terpenoids and steroids](#) [Fructose and mannose metabolism](#)

[Galactose metabolism](#) [Glycolysis / Gluconeogenesis](#) [Insulin signaling pathway](#)

[Metabolic pathways](#) [Starch and sucrose metabolism](#) [Streptomycin biosynthesis](#)

[Type II diabetes mellitus](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.