



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

HLA-C (Human) Recombinant Protein (P02)

Catalog Number: H00003107-P02

Regulation Status: For research use only (RUO)

Product Description: Human HLA-C full-length ORF (AAH02463.1, 1 a.a. - 366 a.a.) recombinant protein with GST tag at N-terminal.

Sequence:

MRVMAPRTLILLLSGALALTETWACSHSMRYFYTAVS
RPGRGEPRIAVGYVDDTQFVRFSDAASPRGEPRAP
WVEQEGPEYWDRETQKYKRQAQTDRLRNLRGYY
NQSEAGSHTLQWMYGCDLGPDRLLRGYDQSAYDG
KDYIALNEDLRSWTAADTAAQITQRKWEAARAAEQQR
AYLEGTVCVEWLRRYLENGKETLQRAEHPKTHVTHHLV
SDHEATLRCWALGFYPAEITLTWQRDGEDQTQDTELV
ETRPAGDGTQKWAAVVPSGEEQRYTCHVQHEGLP
EPLTLRWEPSQPTIPIVGIVAGLAVLAVLAVLGAVVAV
VMCRRKSSGGKGGSCSQAASSNSAQQSDESLIACKA

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 67.2

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3107

Gene Symbol: HLA-C

Gene Alias: D6S204, FLJ27082, HLA-Cw, HLA-Cw12, HLA-JY3, HLC-C, PSORS1

Gene Summary: HLA-C belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by presenting peptides derived from endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the alpha1 and alpha2 domain, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region, and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. Over one hundred HLA-C alleles have been described [provided by RefSeq]