



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### HLA-DOB MaxPab rabbit polyclonal antibody (D01)

**Catalog Number:** H00003112-D01

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit polyclonal antibody raised against a full-length human HLA-DOB protein.

**Immunogen:** HLA-DOB (NP\_002111.1, 1 a.a. ~ 273 a.a) full-length human protein.

**Sequence:**

MGSGWVPWVVALLVNLTRLDSSMTQGTDSPEDFVIQ  
AKADCYFTNGTEKVQFVVRFIFNLEEYVRFSDVGMF  
VALTKLQPDAAEQWNSRLDLLERSRQAVDGVCRHNY  
RLGAPFTVGRKVQPEVTYYPERTPLLHQHNLHCSVT  
GFYPGDIKIKWFLNGQEERAGVMSTGPIRNGDWTFTQ  
VVMLEMTPELGHVYTCLVDHSSLLSPVSVEWRAQSEY  
SWRKMLSGIAAFLGLIFLLVGIQQLRAQKGYVRTQMS  
GNEVSRAVLLPQSC

**Host:** Rabbit

**Reactivity:** Human

**Applications:** IP, WB-Ti, WB-Tr

(See our web site product page for detailed applications information)

**Protocols:** See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Storage Buffer:** No additive

**Storage Instruction:** Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 3112

**Gene Symbol:** HLA-DOB

**Gene Alias:** DOB

**Gene Summary:** HLA-DOB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DOA) and a beta chain (DOB), both anchored in the membrane. It is

located in intracellular vesicles. DO suppresses peptide loading of MHC class II molecules by inhibiting HLA-DM. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. [provided by RefSeq]