



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

HLA-DRB3 (Human) Recombinant Protein (P01)

Catalog Number: H00003125-P01

Regulation Status: For research use only (RUO)

Product Description: Human HLA-DRB3 full-length ORF (AAH08987.1, 1 a.a. - 266 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MVCLKLPGGSSLAALTVTLMLVSSRLAFAGDTRPRFLE
LRKSECHFFNGTERVRYLDRYFHNQEEFLRFDSDVGE
YRAVTELGPRVAESWNSQKDLLEQKRGRVDNYCRHN
YGVGESFTVQRRVHPQVTVPYPAKTQPLQHNNLLVCSV
SGFYPGSIEVRWFRNGQEEKAGVVSTGLIQNGDWTF
QTLVMLETVPRSGEVYTCQVEHPSVTSALTVEWRARS
ESAQSKMLSGVGGFVLGLLFLGAGLFIYFRNQKKGHS
LQPTGFLS

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 55

Applications: AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3125

Gene Symbol: HLA-DRB3

Gene Alias: HLA-DR3B, MGC117330

Gene Summary: HLA-DRB3 belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. DRB1 is expressed at a level five times higher than its paralogues DRB3, DRB4 and DRB5. The presence of DRB3 is linked with allelic variants of DRB1, otherwise it is omitted. There are 4 related pseudogenes: DRB2, DRB6, DRB7, DRB8 and DRB9. [provided by RefSeq]