



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### LMO1 (Human) Recombinant Protein (Q02)

although more rarely than the related gene, LMO2 is disrupted. [provided by RefSeq]

**Catalog Number:** H00004004-Q02

**Regulation Status:** For research use only (RUO)

**Product Description:** Human LMO1 partial ORF (NP\_002306.1, 15 a.a. - 156 a.a.) recombinant protein with GST tag at N-terminal.

**Sequence:**

VQPKGKQKGCAGCNRKIKDRYLLKALDKYWHEDECLK  
CACDCRLGEGSTLYTKANLILCRRDYLRFLFGTTGN  
CAACKLIPAFEMVMRARDNVYHLDCFACQLCNQRFC  
VGDKFFLKNNMILCQMDYEEGQLNGTFESQVQ

**Host:** Wheat Germ (in vitro)

**Theoretical MW (kDa):** 41.25

**Applications:** AP, Array, ELISA, WB-Re  
(See our web site product page for detailed applications information)

**Protocols:** See our web site at  
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Preparation Method:** [in vitro wheat germ expression system](#)

**Purification:** Glutathione Sepharose 4 Fast Flow

**Storage Buffer:** 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

**Storage Instruction:** Store at -80°C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 4004

**Gene Symbol:** LMO1

**Gene Alias:** MGC116692, RBTN1, RHOM1, TTG1

**Gene Summary:** LMO1 encodes a cysteine-rich, two LIM domain transcriptional regulator. It is mapped to an area of consistent chromosomal translocation in chromosome 11, disrupting it in T-cell leukemia,