

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







SMAD1 monoclonal antibody (M05), clone 1C7

Catalog #: H00004086-M05 規格:[100 ug]

List All



Page 1 of 3 2016/5/21




SMAD1 monoclonal antibody (M05), clone 1C7 Western Blot analysis of SMAD1 expression in Hela S3 NE (Cat # L013V3).



Western Blot (Recombinant protein)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)



enlarge this image

Immunoperoxidase of monoclonal antibody to SMAD1 on formalin-fixed paraffin-embedded human lymph node. [antibody concentration 3 ug/ml]



Sandwich ELISA (Recombinant protein)

Detection limit for recombinant GST tagged SMAD1 is approximately 3ng/ml as a capture antibody.



ELISA

Gene Information

Entrez GenelD: 4086

GeneBank NM_005900

Accession#:

Protein <u>NP_005891</u>

Accession#:

Gene Name: SMAD1

Gene Alias: BSP1,JV4-1,JV41,MADH1,MADR1

Gene SMAD family member 1

Description:

Omim ID: <u>601595</u>

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene belongs to the SMAD, a family of

Page 2 of 3 2016/5/21

proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signals of the bone morphogenetic proteins (BMPs), which are involved in a range of biological activities including cell growth, apoptosis, morphogenesis, development and immune responses. In response to BMP ligands, this protein can be phosphorylated and activated by the BMP receptor kinase. The phosphorylated form of this protein forms a complex with SMAD4, which is important for its function in the transcription regulation. This protein is a target for SMAD-specific E3 ubiquitin ligases, such as SMURF1 and SMURF2, and undergoes ubiquitination and proteasome-mediated degradation. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq

Other Designations:

MAD, mothers against decapentaplegic homolog 1,Mad-related protein 1,SMAD, mothers against DPP homolog 1,Sma- and Mad-related protein 1,TGF-beta signaling protein 1,mothers against DPP homolog 1,transforming growth factor-beta signaling protein 1

Gene Pathway

TGF-beta signaling pathway

Related Disease

Cleft Lip Cleft Palate Diabetes Mellitus, Type 1 Diabetic Nephropathies

Genetic Predisposition to Disease Head and Neck Neoplasms Hemochromatosis
Hypertension, Pulmonary Kidney Failure, Chronic Neoplasm Recurrence, Local
Neoplasms, Second Primary Obesity Ovarian Failure, Premature
Polycystic Ovary Syndrome Puberty, Delayed Puberty, Precocious Thrombophilia
Tobacco Use Disorder

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/21