

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in







SMAD2 monoclonal antibody (M09), clone 3B4

Catalog #: H00004087-M09 規格:[100 ug]

List All



Page 1 of 3 2016/5/21



SMAD2 monoclonal antibody (M09), clone 3B4 Western Blot analysis of SMAD2 expression in Hela S3 NE (Cat # L013V3).

Protocol Download

Western Blot (Recombinant protein)

Protocol Download

ELISA

Gene Information

Entrez GenelD: 4087

GeneBank

NM 005901

Accession#:

Protein Accession#: NP 005892

Gene Name:

SMAD2

Gene Alias:

JV18,JV18-1,MADH2,MADR2,MGC22139,MGC34440,hMAD-2,hSMAD2

Gene

SMAD family member 2

Description:

Omim ID:

601366

Gene Ontology: Hyperlink

Gene Summary: The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants encoding the same protein have been observed. [provided by RefSeq

Other Designations: MAD, mothers against decapentaplegic homolog 2, Mad protein homolog, Mad, mothers against decapentaplegic homolog 2, Mad-related protein 2,SMAD, mothers against DPP homolog 2,Sma- and Madrelated protein 2, mother against DPP homolog 2

> Page 2 of 3 2016/5/21

Gene Pathway

Adherens junction Cell cycle Colorectal cancer Pancreatic cancer Pathways in cancer TGF-beta signaling pathway Wnt signaling pathway

Related Disease

Adenocarcinoma Cleft Lip Cleft Palate Colitis, Ulcerative Colorectal Neoplasms

Crohn's disease Esophageal Neoplasms Genetic Predisposition to Disease

Hypertension, Pulmonary Inflammatory Bowel Diseases Liver Cirrhosis Obesity

Osteoporosis Ovarian Failure, Premature Pancreatic cancer Pancreatic Neoplasms

Polycystic Ovary Syndrome Puberty, Delayed Puberty, Precocious

... see more

服務條款 | 隱私權政策 | 著作及商標 | 網站地圖

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.

Page 3 of 3 2016/5/21