



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MAGEA3 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00004102-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-MAGEA3 full-length

Host: Human

Theoretical MW (kDa): 34.65

Quality Control Testing: Transient overexpression cell lysate was tested with Anti-MAGEA3 antibody ([H00004102-B01](#)) by Western Blots.


Western Blot



Lane 1: MAGEA3 transfected lysate (34.65 KDa)

Lane 2: Non-transfected lysate.

SDS-PAGE Gel



MAGEA3 transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [4102](#)

**GeneBank
Accession#:** [NM_005362.3](#)

**Protein
Accession#:** [NP_005353.1](#)

Gene Name: MAGEA3

Gene Alias: HIP8,HYPD,MAGE3,MAGEA6,MGC14613

**Gene
Description:** melanoma antigen family A, 3

Omim ID: [300174](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene is a member of the MAGEA gene family. The members of this family encode proteins with 50 to 80% sequence identity to each other. The promoters and first exons of the MAGEA genes show considerable variability, suggesting that the existence of this gene family enables the same function to be expressed under different transcriptional controls. The MAGEA genes are clustered at chromosomal location Xq28. They have been implicated in some hereditary disorders, such as dyskeratosis congenita. [provided by RefSeq]

**Other
Designations:** MAGE-3 antigen,OTTHUMP00000024220,antigen MZ2-D,melanoma-associated antigen 3

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.