



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MAGEA9 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00004108-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-MAGEA9 full-length

Host: Human

Theoretical MW (kDa): 34.76


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-MAGEA9 antibody (H00004108-B01) by Western Blots.

SDS-PAGE Gel



MAGEA9 transfected lysate.

Western Blot



Lane 1: MAGEA9 transfected lysate (34.76 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [4108](#)

GeneBank Accession#: [NM_005365.4](#)

Protein Accession#: =

Gene Name: MAGEA9

Gene Alias: MAGE9,MGC8421

Gene Description: melanoma antigen family A, 9

Omim ID: [300342](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene is a member of the MAGEA gene family. The members of this family encode proteins with 50 to 80% sequence identity to each other. The promoters and first exons of the MAGEA genes show considerable variability, suggesting that the existence of this gene family enables the same function to be expressed under different transcriptional controls. The MAGEA genes are clustered at chromosomal location Xq28. They have been implicated in some hereditary disorders, such as dyskeratosis congenita. [provided by RefSeq]

Other Designations: MAGE-9 antigen,OTTHUMP00000024214,melanoma-associated antigen 9