



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

MAGOH (Human) Recombinant Protein (P01)

Catalog Number: H00004116-P01

Regulation Status: For research use only (RUO)

Product Description: Human MAGOH full-length ORF (NP_002361.1, 1 a.a. - 146 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MESDFYLRYVVGHKGKFGHEFLEFEFRPDGKLRVANN
SNYKNDVMIRKEAYVHKSVMEELKRIIDDSEITKEDDAL
WPPDRVGRQELEIVIGDEHISFTTSKIGSLIDVNQSKD
PEGLRVFYLVQDLKCLVFLSLIGLHFKIKPI

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 43.6

Applications: AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 4116

Gene Symbol: MAGOH

Gene Alias: MAGOHA

Gene Summary: Drosophila that have mutations in their mago nashi (grandchildless) gene produce progeny with defects in germline assembly and germline development. This gene encodes the mammalian mago

nashi homolog. In mammals, mRNA expression is not limited to the germ plasm, but is expressed ubiquitously in adult tissues and can be induced by serum stimulation of quiescent fibroblasts. [provided by RefSeq]