



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com



[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MET polyclonal antibody (A01)

Catalog # : H00004233-A01

規格 : [50 uL]


List All

Specification	Application Image
Product Description: Mouse polyclonal antibody raised against a partial recombinant MET.	ELISA
Immunogen: MET (NM_000245, 26 a.a. ~ 125 a.a) partial recombinant protein with GST tag.	
Sequence: CKEALAKSEMNVNMKYQLPNFTAETPIQNVILHEHHIFLGATNYIYVLNEED LQKVAEYKTGPVLEHPDCFPCQDCSSKANLSGGVWVDNINMALVVDY	
Host: Mouse	
Reactivity: Human	
Quality Control Testing: Antibody Reactive Against Recombinant Protein.	
Storage Buffer: 50 % glycerol	
Storage Instruction: Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.	
MSDS:  Download	
Datasheet:  Download	
Applications	
ELISA	
Gene Information	
Entrez GeneID: 4233	
GeneBank Accession#: NM_000245	
Protein Accession#: NM_000245	
Gene Name: MET	
Gene Alias: AUTS9,HGFR,RCCP2,c-Met	
Gene Description: met proto-oncogene (hepatocyte growth factor receptor)	
Omim ID: 114550 , 164860 , 605074 , 611015	
Gene Ontology: Hyperlink	
Gene Summary: The proto-oncogene MET product is the hepatocyte growth factor	

receptor and encodes tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. Various mutations in the MET gene are associated with papillary renal carcinoma. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq

Other HGF receptor,OTTHUMP00000069168,SF receptor,met proto-oncogene,met proto-oncogene tyrosine kinase,oncogene MET,scatter factor receptor

Interactome



Gene Pathway

[Adherens junction](#) [Axon guidance](#) [Colorectal cancer](#) [Cytokine-cytokine receptor interaction](#) [Endocytosis](#) [Epithelial cell signaling in Helicobacter pylori infection](#) [Focal adhesion](#) [Melanoma](#) [Pathways in cancer](#) [Renal cell carcinoma](#)

Related Disease

[Adenocarcinoma](#) [Autistic Disorder](#) [Carcinoma, Adenosquamous](#) [Carcinoma, Large Cell](#) [Carcinoma, Non-Small-Cell Lung](#) [Carcinoma, Squamous Cell](#) [Cell Transformation, Neoplastic](#) [Child Development Disorders, Pervasive](#) [Chronic Disease](#) [Cognition Disorders](#) [Disease Progression](#) [Gastrointestinal Diseases](#) [Genetic Predisposition to Disease](#) [Head and Neck Neoplasms](#) [Hyperparathyroidism, Secondary](#) [Lung Neoplasms](#) [Myopia](#) [Nasal Polyps](#) [Neoplasm Metastasis](#)

... see more

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.