



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MF12 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00004241-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-MF12 full-length

Host: Human

Theoretical MW (kDa): 33.33


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-MF12 antibody (H00004241-B01) by Western Blots.

SDS-PAGE Gel



MF12 transfected lysate.

Western Blot



Lane 1: MF12 transfected lysate (33.33 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [4241](#)

**GeneBank
Accession#:** [NM_033316.2](#)

**Protein
Accession#:** [NP_201573.1](#)

Gene Name: MF12

Gene Alias: CD228,FLJ38863,MAP97,MGC4856,MTF1

**Gene
Description:** antigen p97 (melanoma associated) identified by monoclonal antibodies 133.2 and 96.5

Omim ID: [155750](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a cell-surface glycoprotein found on melanoma cells. The protein shares sequence similarity and iron-binding properties with members of the transferrin superfamily. The importance of the iron binding function has not yet been identified. This gene resides in the same region of chromosome 3 as members of the transferrin superfamily. Alternative splicing results in two transcript variants. [provided by RefSeq]

**Other
Designations:** melanoma-associated antigen p97,melanoma-associated antigen p97, isoform 2,melanotransferrin

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.