



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

NAB2 MaxPab rabbit polyclonal antibody (D01)

Catalog Number: H00004665-D01

Regulatory Status: For research use only (RUO)

Product Description: Rabbit polyclonal antibody raised against a full-length human NAB2 protein.

Immunogen: NAB2 (AAH07756.1, 1 a.a. ~ 241 a.a) full-length human protein.

Sequence:

MHRAPSPTAEQPPGGGDSARRTLQPRLKPSARAMAL
PRTLGLQLYRVLQRANLLSYYETFIQQGGDDVQQLC
EAGEEEFLEIMALVGMATKPLHVRRLQKALREWATNP
GLFSQPVPVAVPVSSIPLFKISETAGTRKGSMSNGHGSP
GEKAGSARSFSPKSPLELGEKLSPLPGGPGAGDPRIW
PGRSTPESDVGAGGEEEEAGSRWDWGWSRPTGARD
GTPACLGRVWMDICRLWGHVQG

Host: Rabbit

Reactivity: Human

Applications: IP, WB-Tr

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Storage Buffer: No additive

Storage Instruction: Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 4665

Gene Symbol: NAB2

Gene Alias: MADER, MGC75085

Gene Summary: This gene encodes a member of the family of NGFI-A binding (NAB) proteins, which function in the nucleus to repress transcription induced by some members of the EGR (early growth response) family of transactivators. NAB proteins can homo- or

hetero-multimerize with other EGR or NAB proteins through a conserved N-terminal domain, and repress transcription through two partially redundant C-terminal domains. Transcriptional repression by the encoded protein is mediated in part by interactions with the nucleosome remodeling and deacetylase (NuRD) complex. Alternatively spliced transcript variants have been described, but their biological validity has not been determined. [provided by RefSeq]