



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NFATC3 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00004775-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-NFATC3 full-length

Host: Human

Theoretical MW (kDa): 115.2


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-NFATC3 antibody ([H00004775-B01](#)) by Western Blots.

SDS-PAGE Gel



NFATC3 transfected lysate.

Western Blot



Lane 1: NFATC3 transfected lysate (115.2 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [4775](#)

GeneBank Accession#: [NM_004555.2](#)

Protein Accession#: =

Gene Name: NFATC3

Gene Alias: NFAT4,NFATX

Gene Description: nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 3

Omim ID: [602698](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The product of this gene is a member of the nuclear factors of activated T cells DNA-binding transcription complex. This complex consists of at least two components: a preexisting cytosolic component that translocates to the nucleus upon T cell receptor (TCR) stimulation and an inducible nuclear component. Other members of this family participate to form this complex also. The product of this gene plays a role in the regulation of gene expression in T cells and immature thymocytes. Four transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq]

Other Designations: T cell transcription factor NFAT4,nuclear factor of activated T-cells c3 isoform IE-Xa,nuclear factor of activated T-cells, cytoplasmic 3

Gene Pathway

[Axon guidance](#) [B cell receptor signaling pathway](#) [Natural killer cell mediated cytotoxicity](#) [T cell receptor signaling pathway](#) [VEGF signaling pathway](#) [Wnt signaling pathway](#)

Related Disease

[Cardiovascular Diseases](#) [Diabetes Mellitus, Type 2](#) [Edema](#) [Kidney Failure, Chronic](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.