



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NFYC 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00004802-T01

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-NFYC full-length

Host: Human

Theoretical MW (kDa): 37.2


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-NFYC antibody (H00004802-D01P) by Western Blots.

SDS-PAGE Gel



NFYC transfected lysate.

Western Blot



Lane 1: NFYC transfected lysate (37.20 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [4802](#)

GeneBank [NM_014223.2](#)
Accession#:

Protein [NP_055038.2](#)
Accession#:

Gene Name: NFYC

Gene Alias: CBF-C,CBFC,DKFZp667G242,FLJ45775,H1TF2A,HAP5,HSM,NF-YC

Gene Description: nuclear transcription factor Y, gamma

Omim ID: [605344](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes one subunit of a trimeric complex forming a highly conserved transcription factor that binds with high specificity to CCAAT motifs in the promoters of a variety of genes. The encoded protein, subunit C, forms a tight dimer with the B subunit, a prerequisite for subunit A association. The resulting trimer binds to DNA with high specificity and affinity. Subunits B and C each contain a histone-like motif. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations: CCAAT binding factor subunit C,CCAAT transcription binding factor subunit gamma,OTTHUMP00000009207,OTTHUMP00000009208,OTTHUMP00000009211,OTTHUMP00000009212,histone H1 transcription factor large subunit 2A,transactivator HSM-1,transcription factor NF-Y, C

Gene Pathway

[Antigen processing and presentation](#)

Related Disease

[Carcinoma, Renal Cell](#) [Genetic Predisposition to Disease](#) [Kidney Neoplasms](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.