



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

OAS1 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00004938-T03

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-OAS1 full-length

Host: Human

Theoretical MW (kDa): 40.04


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-OAS1 antibody (H00004938-D03P) by Western Blots.

SDS-PAGE Gel



OAS1 transfected lysate.

Western Blot



Lane 1: OAS1 transfected lysate (40.04 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [4938](#)

GeneBank Accession#: [BC000562.1](#)

Protein Accession#: [AAH00562.1](#)

Gene Name: OAS1

Gene Alias: IFI-4,OIAS,OIASI

Gene Description: 2',5'-oligoadenylate synthetase 1, 40/46kDa

Omim ID: [164350](#), [222100](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes a member of the 2-5A synthetase family, essential proteins involved in the innate immune response to viral infection. The encoded protein is induced by interferons and uses adenosine triphosphate in 2'-specific nucleotidyl transfer reactions to synthesize 2',5'-oligoadenylates (2-5As). These molecules activate latent RNase L, which results in viral RNA degradation and the inhibition of viral replication. The three known members of this gene family are located in a cluster on chromosome 12. Mutations in this gene have been associated with host susceptibility to viral infection. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq]

Other Designations: (2'-5') oligoadenylate synthetase 1,2',5'-oligo A synthetase 1,2',5'-oligoadenylate synthetase 1,2',5'-oligoadenylate synthetase 1 (40-46 kD),2'-5' oligoadenylate synthetase 1 p48 isoform,2'-5' oligoadenylate synthetase 1 p52 isoform,2'-5'-oligoisoadenyla

Related Disease

[Diabetes Mellitus, Type 1](#) [Disease Progression Genetic Predisposition to Disease](#)
[Hepatitis B](#) [Hepatitis B, Chronic](#) [Hepatitis C](#) [Hepatitis C, Chronic](#) [Liver Cirrhosis](#)
[Multiple Sclerosis](#) [Multiple Sclerosis, Relapsing-Remitting](#) [Ovarian Neoplasms](#)
[Severe Acute Respiratory Syndrome](#) [West Nile Fever](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.