



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

REG3A 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00005068-T03

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-REG3A full-length

Host: Human

Theoretical MW (kDa): 19.4


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-REG3A antibody ([H00005068-D01P](#)) by Western Blots.

SDS-PAGE Gel



REG3A transfected lysate.

Western Blot



Lane 1: REG3A transfected lysate (19.40 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS: [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [5068](#)

GeneBank Accession#: [NM_002580](#)

Protein Accession#: [NP_002571.1](#)

Gene Name: REG3A

Gene Alias: HIP,INGAP,PAP,PAP-H,PAP1,PBCGF,REG-III,REG3

Gene Description: regenerating islet-derived 3 alpha

Omim ID: [167805](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene encodes a pancreatic secretory protein that may be involved in cell proliferation or differentiation. It has similarity to the C-type lectin superfamily. The enhanced expression of this gene is observed during pancreatic inflammation and liver carcinogenesis. Multiple alternatively spliced transcript variants encoding the same protein have been described for this gene but the full length nature of some transcripts is not yet known. [provided by RefSeq]

Other Designations: PAP homologous protein,hepatocarcinoma-intestine-pancreas,pancreatic beta cell growth factor,pancreatitis-associated protein,proliferation-inducing protein 34,proliferation-inducing protein 42

Related Disease

[Alzheimer Disease](#) [Alzheimer disease](#) [Cardiovascular Diseases](#) [Diabetes Complications](#)
[Metabolic Syndrome X](#) [Neoplasms](#) [Osteoporosis](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.