



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

MAPK13 (Human) Recombinant Protein (Q01)

Catalog Number: H00005603-Q01

Regulation Status: For research use only (RUO)

Product Description: Human MAPK13 partial ORF (AAH00433, 251 a.a. - 365 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

NDKAAKSYIQSLPQTTPRKDFTQLFPRASPQAADLLEK
MLELDVDRRLTAAQALHPFFEPFRDPEEETEAQQPF
DDSLEHEKLTVDEWKQHYYKEIVNFSPARKDSRRRSG
MKL

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 38.28

Applications: AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 5603

Gene Symbol: MAPK13

Gene Alias: MGC99536, PRKM13, SAPK4, p38delta

Gene Summary: The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such

as proliferation, differentiation, transcription regulation and development. This kinase is closely related to p38 MAP kinase, both of which can be activated by proinflammatory cytokines and cellular stress. MAP kinase kinases 3, and 6 can phosphorylate and activate this kinase. Transcription factor ATF2, and microtubule dynamics regulator stathmin have been shown to be the substrates of this kinase. [provided by RefSeq]