



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

PSMA3 (Human) Recombinant Protein (P01)

Catalog Number: H00005684-P01

Regulation Status: For research use only (RUO)

Product Description: Human PSMA3 full-length ORF (AAH05265, 1 a.a. - 248 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

```
MSSIGTGYDLSASTFSPDGRVVFQVEYAMKA VENSSTAI  
GIRCKDGVVFGVEKLVLSKLYEEGSNKRLFNVDRHVG  
MAVAGLLADARSLADIAREEASNFRSNFGYNIPLKHLA  
DRVAMYVHAYTLYSAVRPFGCSVNDGAQLYMIDPSGV  
SYGYWGC AIGKARQAAKTEIEKLQMKEMTCRDIVKEV  
AKIYIVHDEVKDKAFELELSWVGELTNGRHEIVPKDIR  
EEAEKYAKESLKEEDESDDDNM
```

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 53.02

Applications: AP, Array, ELISA, WB-Re

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 5684

Gene Symbol: PSMA3

Gene Alias: HC8, MGC12306, MGC32631, PSC3

Gene Summary: The proteasome is a multicatalytic

proteasome complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. Two alternative transcripts encoding different isoforms have been identified. [provided by RefSeq]