



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

PSMB7 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00005695-T02

規格 : [100 uL]

[List All](#)

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-PSMB7 full-length

Host: Human

Theoretical MW (kDa): 30.58


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-PSMB7 antibody (H00005695-B02) by Western Blots.

SDS-PAGE Gel



PSMB7 transfected lysate.

Western Blot



Lane 1: PSMB7 transfected lysate (30.58 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS: [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [5695](#)

GeneBank [BC000509](#)
Accession#:

Protein [AAH00509](#)
Accession#:

Gene Name: PSMB7

Gene Alias: Z

Gene Description: proteasome (prosome, macropain) subunit, beta type, 7

Omim ID: [604030](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit in the proteasome. Expression of this catalytic subunit is downregulated by gamma interferon and proteolytic processing is required to generate a mature subunit. This subunit is not present in the immunoproteasome and is replaced by catalytic subunit 2i (proteasome beta 10 subunit). [provided by RefSeq]

Other Designations: OTTHUMP00000022798,macropain chain Z,multicatalytic endopeptidase complex chain Z,proteasome beta 7 subunit,proteasome catalytic subunit 2,proteasome subunit Z,proteasome subunit alpha,proteasome subunit beta 7

Gene Pathway

[Proteasome](#)

Related Disease

[Kidney Failure, Chronic Tobacco Use Disorder](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.