



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

PSMD4 (Human) Recombinant Protein (P01)

Catalog Number: H00005710-P01

Regulation Status: For research use only (RUO)

Product Description: Human PSMD4 full-length ORF (NP_002801.1, 1 a.a. - 377 a.a.) recombinant protein with GST-tag at N-terminal.

Sequence:

MVLESTMVCVDNSEYMRNGDFLPTRLQAQQDAVNIV
CHSKTRSNPENNVGLITLANDCEVLTTLPDTGRILSKL
HTVQPKGKITFCTGIRVAHLALKHRQGNHMKMRIAIFV
GSPVEDNEKDLVKLAKRLKKEKVNVDIINFGEEEVNTE
KLTAfvNTLNGKDGtGSHLVTVPPGpSLADALISSPILA
GEGGAMLGLGASDFEFGVDPSADPELALALRVSMEE
QRQRQEEEEARRAAAASAAEAGIATTGTEDSDDALLKM
TISQQEFGRtGLPDLSSMTEEEQIAYAMQMSLQGAEF
GQAESADIDASSAMDTSEPAKEEDDYDVMQDPEFLQ
SVLENLPGVDPNNEAIRNAMGSLASQATKDGKDKKKE
EDKK

Host: Wheat Germ (in vitro)

Theoretical MW (kDa): 67.1

Applications: AP, Array, ELISA, WB-Re
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Preparation Method: [in vitro wheat germ expression system](#)

Purification: Glutathione Sepharose 4 Fast Flow

Storage Buffer: 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 5710

Gene Symbol: PSMD4

Gene Alias: AF, AF-1, ASF, MCB1, Rpn10, S5A, pUB-R5

Gene Summary: The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the non-ATPase subunits of the 19S regulator lid. Pseudogenes have been identified on chromosomes 10 and 21. [provided by RefSeq]