



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## PSME2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00005721-T01

規格 : [ 100 uL ]

List All

### Specification

**Transfected Cell Line:** 293T


**Plasmid:** pCMV-PSME2 full-length

**Host:** Human

**Theoretical MW (kDa):** 26.4


**Quality Control Testing:** Transient overexpression cell lysate was tested with Anti-PSME2 antibody (H00005721-B01) by Western Blots.

#### SDS-PAGE Gel



PSME2 transfected lysate.

#### Western Blot



Lane 1: PSME2 transfected lysate ( 26.4 KDa)

Lane 2: Non-transfected lysate.

**Storage Buffer:** 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

**Storage Instruction:** Store at -80°C. Aliquot to avoid repeated freezing and thawing.

**MSDS:**  [Download](#)

### Applications

### Application Image

Western Blot

## Western Blot

### Gene Information

Entrez GeneID: [5721](#)

GeneBank [BC004368.1](#)  
Accession#:

Protein =  
Accession#:

Gene Name: PSME2

Gene Alias: PA28B,PA28beta,REGbeta

Gene Description: proteasome (prosome, macropain) activator subunit 2 (PA28 beta)

Omim ID: [602161](#)

Gene Ontology: [Hyperlink](#)

**Gene Summary:** The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. The immunoproteasome contains an alternate regulator, referred to as the 11S regulator or PA28, that replaces the 19S regulator. Three subunits (alpha, beta and gamma) of the 11S regulator have been identified. This gene encodes the beta subunit of the 11S regulator, one of the two 11S subunits that is induced by gamma-interferon. Three beta and three alpha subunits combine to form a heterohexameric ring. Six pseudogenes have been identified on chromosomes 4, 5, 8, 10 and 13. [provided by RefSeq]

**Other Designations:** 11S regulator complex beta subunit,MCP activator, 31-kD subunit,activator of multicatalytic protease subunit 2,cell migration-inducing protein 22,proteasome activator 28-beta,proteasome activator hPA28 subunit beta,proteasome activator subunit 2

### Gene Pathway

[Antigen processing and presentation](#) [Proteasome](#)

### Related Disease

[Disease Progression](#) [Disease Susceptibility](#) [HIV Infections](#)