



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PTPN7 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00005778-T02

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-PTPN7 full-length

Host: Human

Theoretical MW (kDa): 39.71


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-PTPN7 antibody (H00005778-B02) by Western Blots.

SDS-PAGE Gel



PTPN7 transfected lysate.

Western Blot



Lane 1: PTPN7 transfected lysate (39.71 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [5778](#)

GeneBank Accession#: [NM_002832.2](#)

Protein Accession#: [NP_002823.2](#)

Gene Name: PTPN7

Gene Alias: BPTP-4,HEPTP,LC-PTP,LPTP,PTPNI

Gene Description: protein tyrosine phosphatase, non-receptor type 7

Omim ID: [176889](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This gene is preferentially expressed in a variety of hematopoietic cells, and is an early response gene in lymphokine stimulated cells. The noncatalytic N-terminus of this PTP can interact with MAP kinases and suppress the MAP kinase activities. This PTP was shown to be involved in the regulation of T cell antigen receptor (TCR) signaling, which was thought to function through dephosphorylating the molecules related to MAP kinase pathway. Two alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations: OTTHUMP00000034115,dual specificity phosphatase 1,hematopoietic protein-tyrosine phosphatase,protein-tyrosine phosphatase, nonreceptor-type, stress induced

Gene Pathway

[MAPK signaling pathway](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.