



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

RRM2 MaxPab rabbit polyclonal antibody (D01)

Catalog Number: H00006241-D01

Regulatory Status: For research use only (RUO)

Product Description: Rabbit polyclonal antibody raised against a full-length human RRM2 protein.

Immunogen: RRM2 (NP_001025.1, 1 a.a. ~ 389 a.a) full-length human protein.

Sequence:

```
MLSLRVPLAPITDPQQLQLSPLKGLSLVDKENTPPALS  
GTRVLASKTARRIFQEPTTEPKTKAAAPGVEDEPLLREN  
PRRFVIFPIEYHDIWQMYKKAASFMTAAEVDLSKDIQ  
HWESLKPEERYFISHVLAFFAASDGIVNENLVERFSQE  
VQITEARCFYGFQIAMENIHSEMYSLIDITYIKDKERE  
FLFNAIETMPCVKKKADWALRWIGDKEATYGERVAVF  
AAVEGIFFSGSFASIFWLKKRGLMPGLTFSNELISRDE  
GLHCDFACLMTFKHLVHKPSEERVREIINAVRIEQEFLT  
EALPVKLIQMNCTLMKQYIEFVADRLMLELGFVKVFRV  
ENPFDFMENISLEGKTNFFEKRVGEYQRMGMSSPTE  
NSFTLDADF
```

Host: Rabbit

Reactivity: Human

Applications: IP, WB-Ti, WB-Tr

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Storage Buffer: No additive

Storage Instruction: Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6241

Gene Symbol: RRM2

Gene Alias: R2, RR2M

Gene Summary: This gene encodes one of two

non-identical subunits for ribonucleotide reductase. This reductase catalyzes the formation of deoxyribonucleotides from ribonucleotides. Synthesis of the encoded protein (M2) is regulated in a cell-cycle dependent fashion. Transcription from this gene can initiate from alternative promoters, which results in two isoforms that differ in the lengths of their N-termini. Related pseudogenes have been identified on chromosomes 1 and X. [provided by RefSeq]