



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

SNAI1 MaxPab rabbit polyclonal antibody (D01)

Catalog Number: H00006615-D01

Regulatory Status: For research use only (RUO)

Product Description: Rabbit polyclonal antibody raised against a full-length human SNAI1 protein.

Immunogen: SNAI1 (NP_005976.2, 1 a.a. ~ 264 a.a) full-length human protein.

Sequence:

MPRSFLVRKPSDPNRKPNYSELQDSNPEFTFQQPYD
QAHLLAAIPPEILNPTASLPMLIWDSVLAPQAQPIAWA
SLRLQESPRVAELTSLSDEDSGKGSQPPSPSPAPSS
FSSTSVSLEAEAYAAFPGLGQVPKQLAQLSEAKDLQ
ARKAFNCKYCNKEYLSLGALKMHIRSHTLPCVCGTTCG
KAFSRPWLLQGHVRTHTGKPFSCPHCSRAFADRSN
LRAHLQTHSDVKKYQCACARTFSRMSLLHKKHYESG
CSGCPR

Host: Rabbit

Reactivity: Human

Applications: IP, WB-Tr

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Storage Buffer: No additive

Storage Instruction: Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6615

Gene Symbol: SNAI1

Gene Alias: SLUGH2, SNA, SNAH, dJ710H13.1

Gene Summary: The Drosophila embryonic protein snail is a zinc finger transcriptional repressor which downregulates the expression of ectodermal genes within the mesoderm. The nuclear protein encoded by

this gene is structurally similar to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing embryo. At least two variants of a similar processed pseudogene have been found on chromosome 2. [provided by RefSeq]