



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

STAT6 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00006778-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-STAT6 full-length

Host: Human

Theoretical MW (kDa): 93.28


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-STAT6 antibody (H00006778-B01) by Western Blots.

SDS-PAGE Gel



STAT6 transfected lysate.

Western Blot



Lane 1: STAT6 transfected lysate (93.28 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Western Blot

Gene Information

Entrez GeneID: [6778](#)

GeneBank Accession#: [NM_003153.3](#)

Protein Accession#: [NP_003144.3](#)

Gene Name: STAT6

Gene Alias: D12S1644,IL-4-STAT,STAT6B,STAT6C

Gene Description: signal transducer and activator of transcription 6, interleukin-4 induced

Omim ID: [601512](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4. Knockout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expression of cell surface markers, and class switch of immunoglobulins. [provided by RefSeq]

Other Designations: STAT, interleukin4-induced,signal transducer and activator of transcription 6,transcription factor IL-4 STAT

Gene Pathway

[Jak-STAT signaling pathway](#)

Related Disease

[Anaphylaxis](#) [Arthritis](#) [Rheumatoid](#) [Ascariasis](#) [Asthma](#) [Asthma](#) [Birth Weight](#) [Breast cancer](#) [Breast Neoplasms](#) [Bronchial Hyperreactivity](#) [Bronchiolitis](#) [Viral](#) [Cardiovascular Diseases](#) [Colitis](#) [Ulcerative](#) [Confusion](#) [Crohn Disease](#) [Crohn's disease](#) [Dermatitis](#) [Atopic](#) [Diabetes Mellitus](#) [Type 2 Disease Progression](#) [Drug Hypersensitivity](#)

... see more