



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TANK 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00010010-T02

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-TANK full-length

Host: Human

Theoretical MW (kDa): 13.2


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-TANK antibody (H00010010-B02) by Western Blots.

SDS-PAGE Gel



TANK transfected lysate.

Western Blot



Lane 1: TANK transfected lysate (13.2 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [10010](#)

GeneBank [NM_133484.1](#)
Accession#:

Protein =
Accession#:

Gene Name: TANK

Gene Alias: I-TRAF, TRAF2

Gene Description: TRAF family member-associated NFkB activator

Omim ID: [603893](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: The TRAF (tumor necrosis factor receptor-associated factor) family of proteins associate with and transduce signals from members of the tumor necrosis factor receptor superfamily. The protein encoded by this gene is found in the cytoplasm and can bind to TRAF1, TRAF2, or TRAF3, thereby inhibiting TRAF function by sequestering the TRAFs in a latent state in the cytoplasm. For example, the protein encoded by this gene can block TRAF2 binding to LMP1, the Epstein-Barr virus transforming protein, and inhibit LMP1-mediated NF-kappa-B activation. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations: TRAF family member-associated NF-KAPPA-B activator, TRAF interacting protein TANK

Related Disease

[Arthritis](#), [Rheumatoid Cardiovascular Diseases](#), [Diabetes Mellitus, Type 2](#), [Edema](#), [Genetic Predisposition to Disease](#), [Hematologic Diseases](#), [Hepatitis C](#), [Hodgkin Disease](#), [Lymphoproliferative Disorders](#), [Multiple Myeloma](#), [Occupational Diseases](#), [Osteoporosis](#), [Postmenopausal](#), [Waldenstrom Macroglobulinemia](#), [Werner syndrome](#)

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.