



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TNK2 MaxPab rabbit polyclonal antibody (D01)

Catalog Number: H00010188-D01

Regulatory Status: For research use only (RUO)

Product Description: Rabbit polyclonal antibody raised against a full-length human TNK2 protein.

Immunogen: TNK2 (AAH08884.1, 1 a.a. ~ 352 a.a) full-length human protein.

Sequence:

MQPEEGTGWLLELLSEVQLQQYFLRLRDDLNVTRL
SHFEYVKNEDEKIGMGRPGQRRLEAVKRRKALCKRK
SWMSKVFSGKRLEAEFPPHHSQSTFRKTSPAPGGPA
GEGPLQSLTCLIGEKDLRLLLEKLGDSFVVRRGEWD
APSGKTVSVAVKCLKPDVLSQPEAMDDFIREVNAMHS
LDHRNLIRLYGVVLTTPMKMVTTELAPLGSLDRLRKHQ
GHFLLGTLSTRYAVQVAEGMGYLESKRFIHRDLAARNLL
LATRDLVKIGDFGLMRALPQNDDHYVMQEHRKVPFA
WCAPESLKPPWRDISASSSTQFPHAVPCFPTSLAKLL
LRHSVPASSREIKLVSILC

Host: Rabbit

Reactivity: Human

Applications: IF, IP, WB-Ce, WB-Tr

(See our web site product page for detailed applications information)

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Storage Buffer: No additive

Storage Instruction: Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 10188

Gene Symbol: TNK2

Gene Alias: ACK, ACK1, FLJ44758, FLJ45547, p21cdc42Hs

Gene Summary: This gene encodes a tyrosine kinase

that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length nature of only two transcript variants has been determined. [provided by RefSeq]