



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### RAMP2 (Human) Recombinant Protein (P01)

**Catalog Number:** H00010266-P01

**Regulation Status:** For research use only (RUO)

**Product Description:** Human RAMP2 full-length ORF ( NP\_005845.2, 1 a.a. - 175 a.a.) recombinant protein with GST-tag at N-terminal.

**Sequence:**

MASLRVERAGGPRLPRTRVGRPAALRLLLLLGAVLNP  
HEALAQPLPTTGTPGSEGGTVKNYETAVQFCWNHYK  
DQMDPIEKDWCDWAMISRPYSTLRDCLFHFAELFDLG  
FPNPLAERIIIFETHQIHFANCSLVQPTFSDPPEDVLLAMI  
IAPICLIPFLITLVVWRKSEAQA

**Host:** Wheat Germ (in vitro)

**Theoretical MW (kDa):** 46

**Applications:** AP, Array, ELISA, WB-Re  
(See our web site product page for detailed applications information)

**Protocols:** See our web site at  
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Preparation Method:** [in vitro wheat germ expression system](#)

**Purification:** Glutathione Sepharose 4 Fast Flow

**Storage Buffer:** 50 mM Tris-HCl, 10 mM reduced Glutathione, pH=8.0 in the elution buffer.

**Storage Instruction:** Store at -80°C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 10266

**Gene Symbol:** RAMP2

**Gene Alias:** -

**Gene Summary:** The protein encoded by this gene is a member of the RAMP family of single-transmembrane-domain proteins, called receptor

(calcitonin) activity modifying proteins (RAMPs). RAMPs are type I transmembrane proteins with an extracellular N terminus and a cytoplasmic C terminus. RAMPs are required to transport calcitonin-receptor-like receptor (CRLR) to the plasma membrane. CRLR, a receptor with seven transmembrane domains, can function as either a calcitonin-gene-related peptide (CGRP) receptor or an adrenomedullin receptor, depending on which members of the RAMP family are expressed. In the presence of this (RAMP2) protein, CRLR functions as an adrenomedullin receptor. The RAMP2 protein is involved in core glycosylation and transportation of adrenomedullin receptor to the cell surface. [provided by RefSeq]