



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LILRB2 293T Cell Transient Overexpression Lysate(Denatured)

Catalog # : H00010288-T01

規格 : [100 uL]

List All

Specification

Transfected Cell Line: 293T


Plasmid: pCMV-LILRB2 full-length

Host: Human

Theoretical MW (kDa): 21.2


Quality Control Testing: Transient overexpression cell lysate was tested with Anti-LILRB2 antibody (H00010288-B01) by Western Blots.

SDS-PAGE Gel



LILRB2 transfected lysate.

Western Blot



Lane 1: LILRB2 transfected lysate (21.2 KDa)

Lane 2: Non-transfected lysate.

Storage Buffer: 1X Sample Buffer (50 mM Tris-HCl, 2% SDS, 10% glycerol, 300 mM 2-mercaptoethanol, 0.01% Bromophenol blue)

Storage Instruction: Store at -80°C. Aliquot to avoid repeated freezing and thawing.

MSDS:  [Download](#)

Applications

Application Image

Western Blot

Western Blot

Gene Information

Entrez GeneID: [10288](#)

GeneBank Accession#: [BC041708.1](#)

Protein Accession#: =

Gene Name: LILRB2

Gene Alias: CD85D,ILT4,LILRA6,LIR-2,LIR2,MIR-10,MIR10

Gene Description: leukocyte immunoglobulin-like receptor, subfamily B (with TM and ITIM domains), member 2

Omim ID: [604815](#)

Gene Ontology: [Hyperlink](#)

Gene Summary: This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]

Other Designations: Ig-like transcript 4,OTTHUMP00000067358,OTTHUMP00000067463,eukocyte immunoglobulin-like receptor, subfamily A (with TM domain), member 6,immunoglobulin-like transcript 4,leukocyte immunoglobulin-like receptor 2,leukocyte immunoglobulin-like receptor subfa

[服務條款](#) | [隱私權政策](#) | [著作及商標](#) | [網站地圖](#)

©2016 亞諾法生技股份有限公司 Abnova Corporation. 版權所有.