



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Anti-HOXD12 Antibody (Glucose Oxidase)

Product Number: AC21-2785-13

Overview

Host Species: Goat

Clonality: Polyclonal

Purity: Affinity purified using the immunizing peptide. The affinity purified antibody is then conjugated to glucose oxidase (GOx) which is an enzyme that catalyzes glucose producing hydrogen peroxide.

Conjugate: Glucose Oxidase

Physical Characteristics

Amount: 0.05 mg

Volume: 0.1 ml

Concentration: 0.5 mg/ml

Buffer: 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH 7.3 with 0.5% bovine serum albumin.

Storage: Aliquot and store at -20°C. Minimize freezing and thawing.

Specificity

Tested Species Reactivity: Human

Predicted Species Reactivity: Human

Negative Species Reactivity: Unknown

Immunogen: HOXD12 antibody (Glucose Oxidase) was raised against a synthetic peptide of HOXD12. The HOXD12 peptide immunogen had the following sequence: C-PTAKDGPEEQAK

Blocking Peptide: HOXD12 peptide, catalog number AC21-2785-P, is the immunizing peptide for this product and can be used as a negative control to block antibody (Glucose Oxidase) binding in a competition or adsorption assay.

Target Information

Accession Number(s): NP_067016.3

Gene ID Number(s): 3238 (human)

Alternative Names: Homeobox protein Hox D12; Homeobox protein Hox-4H; Homeobox protein Hox-D12; Hox 4.7, mouse, homolog of; Hox 4H; HOX4H; HOXD12; HXD12_HUMAN



Application Details

Tested Applications: ELISA, WB


ELISA: Detection limit dilution 1:1000

Western Blot: Recommended starting concentration for western blot is 2-3 μ g/ml.

Immunohistochemistry: n/a

Immunocytochemistry: n/a

Product Images



Western blot using HOXD12 antibody (unconjugated) (2 μ g/ml) to detect endogenous HOXD12 in human heart lysate (35 μ g total protein in RIPA buffer).