



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Technically
Speaking

CEDARLANE[®] 
www.cedarlanelabs.com

Conveniently Delivering You Today's Innovations
for the Science of Tomorrow™

Mouse Anti-CD45RO Monoclonal Antibody

CLX545AP
CLX545F
CLX545PE

Clone: UCHL1

Isotype: Mouse IgG2a

Specificity:

The antibody UCHL1 recognizes CD45RO, a 180 kDa low molecular weight isoform of the leukocyte common antigen (LCA). The antigen is expressed on a subset of memory/activated T cells and on cortical thymocytes.

HLDA III; WS Code NL 826

HLDA III; WS Code T 128

HLDA IV; WS Code NL 31

HLDA V; WS Code BP BP460

HLDA V; WS Code T T-081

HLDA V; WS Code T T-CD45.43

Immunogen: Human IL-2 dependent T cells.

Species Reactivity: Human

Application: Flow Cytometry; Immunohistochemistry (paraffin and frozen sections), Immunoprecipitation.

Note: This product does not require protein digestion pre-treatment of paraffin sections. This product does not require antigen retrieval using heat treatment prior to staining of paraffin sections. Positive tissue is human tonsil.

Conjugate Preparation:

The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) and R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

Presentation:

Purified: 0.1 mg (1 mg/mL) purified IgG buffered in PBS with 15 mM sodium azide, approx. pH 7.4. (Purified from hybridoma culture supernatant by protein-G affinity chromatography).

FITC: 2 mL of FITC conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

PE: 2 mL of PE conjugated IgG buffered in stabilizing phosphate buffered saline (PBS) solution containing 15mM sodium azide. Sufficient for 100 tests.

Continued Overleaf.....

Visit our website for your local distributor.

CEDARLANE[®] 

www.cedarlanelabs.com

An ISO 9001:2000 and ISO 13485:2003
registered company.

In CANADA: **Toll Free: 1-800-268-5058**

4410 Paletta Court, Burlington, ON L7L 5R2 ph: (289) 288-0001, fax: (289) 288-0020
e-mail: general@cedarlanelabs.com

In the USA: **Toll Free: 1-800-721-1644**

1210 Turrentine Street, Burlington, NC 27215 ph: (336) 513-5135, fax: (336) 513-5138
e-mail: service@cedarlanelabs.com

Storage / Stability:

Store in the dark at 2-8°C. Do not freeze all formats. Avoid prolonged exposure to light of conjugates. Do not use after expiration date stamped on vial label.

Usage:

Recommended dilutions for Flow Cytometry analysis of human blood cells using:

Purified: 10 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.

FITC: 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.

PE: 20 µl reagent / 100 µl of whole blood or 10⁶ cells in a suspension.

Background:

CD45RO is the shortest isoform of a receptor-type protein tyrosine phosphatase, CD45 glycoprotein. CD45 is crucial in lymphocyte development and antigen signaling, serving as an important regulator of Src-family kinases, promotes cell survival by modulating integrin-mediated signal transduction pathway and is also involved in DNA fragmentation during apoptosis. CD45 isoforms differ in their extracellular domains, whereas they share identical transmembrane and cytoplasmic domains. These isoforms differ in their ability to translocate into the glycosphingolipid-enriched membrane domains and their expression depends on cell type and physiological state of the cell. CD45R0 is expressed e.g. on macrophages, CD8+ T cells, activated T cells and myeloma cells.

References:

- *Li FJ, Tsuyama N, Ishikawa H, Obata M, Abroun S, Liu S, Otsuyama K, Zheng X, Ma Z, Maki Y, Kawano MM: A rapid translocation of CD45RO but not CD45RA to lipid rafts in IL-6-induced proliferation in myeloma. *Blood*. 2005 Apr 15;105(8):3295-302.
- *Cosenza-Nashat MA, Kim MO, Zhao ML, Suh HS, Lee SC: CD45 isoform expression in microglia and inflammatory cells in HIV-1 encephalitis. *Brain Pathol*. 2006 Oct;16(4):256-65.
- *Dawes R, Petrova S, Liu Z, Wraith D, Beverley PC, Tchilian EZ. Combinations of CD45 isoforms are crucial for immune function and disease. *J Immunol*. 2006 Mar 15;176(6):3417-25.
- *Bijian K, Zhang L, Shen SH: Collagen-mediated survival signaling is modulated by CD45 in Jurkat T cells. *Mol Immunol*. 2007 Jul;44(15):3682-90.
- *Desharnais P, Dupéré-Minier G, Hamelin C, Devine P, Bernier J: Involvement of CD45 in DNA fragmentation in apoptosis induced by mitochondrial perturbing agents. *Apoptosis*. 2007 Dec 19
- *Norton AJ, Ramsay AD, Smith SH, Beverley PC, Isaacson PG: Monoclonal antibody (UCHL1) that recognises normal and neoplastic T cells in routinely fixed tissues. *J Clin Pathol*. 1986 Apr;39(4):399-405.
- *Smith SH, Brown MH, Rowe D, Callard RE, Beverley PC: Functional subsets of human helper-inducer cells defined by a new monoclonal antibody, UCHL1. *Immunology*. 1986 May;58(1):63-70.
- *Beverley PC.: Human T cell subsets. *Immunol Lett*. 1987 Apr;14(4):263-7.
- *Leukocyte Typing III., McMichael A. J. et al (Eds.), Oxford University Press (1987).
- *Akbar AN, Terry L, Timms A, Beverley PC, Janossy G: Loss of CD45R and gain of UCHL1 reactivity is a feature of primed T cells. *J Immunol*. 1988 Apr 1;140(7):2171-8.
- *Terry LA, Brown MH, Beverley PC: The monoclonal antibody, UCHL1, recognizes a 180,000 MW component of the human leucocyte-common antigen, CD45. *Immunology*. 1988 Jun;64(2):331-6.
- *Beverley PC, Merckenschlager M, Terry L: Phenotypic diversity of the CD45 antigen and its relationship to function. *Immunol Suppl*. 1988;1:3-5.
- *Merckenschlager M, Terry L, Edwards R, Beverley PC: Limiting dilution analysis of proliferative responses in human lymphocyte populations defined by the monoclonal antibody UCHL1: implications for differential CD45 expression in T cell memory formation. *Eur J Immunol*. 1988 Nov;18(11):1653-61.
- *Leukocyte Typing IV., Knapp W. et al. (Eds.), Oxford University Press (1989).
- *Leukocyte Typing V., Schlossman S. et al. (Eds.), Oxford University Press (1995).
- *And many other.

Laboratory Reagent For Research Use Only