



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com


www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Anti-Mouse Hepsin In Vivo Antibody - Low Endotoxin (2D5-1.9) [ICH1160]

SKU: ICH1160

Link: <https://www.ichor.bio/product/anti-mouse-hepsin-in-vivo-antibody-low-endotoxin-2d5-1-9-ich1160/>



Product Information

Category: anti-mouse, Low Endotoxin, Ultra Low Endotoxin

Size: 1mg, 5mg, 25mg, 50mg, 100mg

Endotoxin Level: Low, Ultra low

Product Description

Product Benefits:

ichorbio's anti-Hepsin In Vivo Antibody - Low Endotoxin (2D5-1.9) is manufactured in a cGMP compliant, ISO Quality Standard 9001:2015 facility. ichorbio's low endotoxin antibodies have half the endotoxin of comparable antibodies from our competitors (<https://www.ichor.bio/comparing-ichorbio-to-bio-x-cell-biolegend/>), at less than 1.0 EU/mg. If ichorbio's low endotoxin antibodies are not low enough we also offer ultra low endotoxin antibodies which have even less endotoxin (<0.75EU/mg) at an even higher purity (98% versus 95%). ichorbio offers Amazon vouchers or donations to the NC3Rs for reviews of this product: click here (<https://www.ichor.bio/amazon-vouchers/>), for more information. ichorbio: the best antibodies for *in vivo* research.

Target:

Hepsin

Clone:

2D5-1.9

Isotype:

Mouse IgG2a κ

Other Names:

Serine protease hepsin, hpn

Uniprot:

P05981 (<https://www.uniprot.org/uniprot/P05981>).

Host:

Mouse

Species Reactivity:

Human

Specificity:

Clone 2D5 recognizes Human hepsin. It binds to a separate epitope than clone 3H10.1. These clones do not interfere with one another.

Purification Method:

This monoclonal antibody was purified using multi-step affinity chromatography methods such as Protein A or G depending on the species and isotype.

Antigen Distribution:

Hepsin is expressed on the surface of epithelial cells including the liver, kidney, prostate, and thyroid in human tissues.

Background:

Hepsin is a type II transmembrane serine protease (TTSP) expressed on the surface of epithelial cells including the liver, kidney, prostate, and thyroid in human tissues. The physiological function of hepsin is unclear, although, In vitro studies have shown that hepsin activates blood clotting factors VII, XII, and IX, pro-

urokinase (pro-uPA), and pro-hepatocyte growth factor (pro-HGF). The over-expression of hepsin has been implicated in several types of cancer, especially ovarian and prostate, which makes it an attractive diagnostic marker for cancers. Most notably, hepsin has been identified as one of the most highly induced genes in prostate cancer, and this over-expression is correlated with the cancer progression and metastasis. Furthermore anti-hepsin antibodies have been shown to inhibit the invasion of human prostate cancer cells.¹

Immunogen:

Human hepsin protein

Concentration:

1.0 mg/ml

Formulation:

This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (PBS) pH 7.2 - 7.4, 150 mM NaCl with no carrier protein, potassium or preservatives added. BSA and azide free

Purity:

>95% by SDS-PAGE and HPLC

>98% by SDS-PAGE and HPLC

Endotoxin:

≤ 1.0 EU/mg as determined by the LAL method

≤ 0.75 EU/mg as determined by the LAL method

Aggregation:

Aggregation level ≤ 5%

Aggregation level ≤ 1%

IMPACT Pathogen Test:

We use the IMPACT test generated by IDEXX Laboratories to guarantee our Ultra Low Endotoxin antibodies are pathogen free. Our mouse antibodies are tested for: Mycoplasma spp. Mycoplasma pulmonis Sendai virus Mouse hepatitis virus Pneumonia virus of mice Minute virus of mice Mouse parvovirus (MPV1-5) Theiler's murine encephalomyelitis virus Murine norovirus Reovirus 3 Mouse rotavirus Ectromelia virus Lymphocytic choriomeningitis virus Polyoma virus Lactate dehydrogenase-elevating virus Mouse adenovirus (MAD1, MAD2) Mouse cytomegalovirus K virus Mouse thymic virus Hantaan virus Corynebacterium bovis Corynebacterium spp. (HAC2)

Storage:

This antibody is stable for at least one week when stored sterile at 2-8°C. For long term storage aseptically aliquot in working volumes without diluting and store at -80°C. Avoid Repeated Freeze Thaw Cycles.

Applications:

ELISA, FC, WB

Application Notes:

Each investigator should determine their own optimal working dilution for specific applications.

Use:

Products are for research use only. Not for use in diagnostic or therapeutic procedures.