



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



872 W. Baltimore Pike, West Grove, PA 19390
800-367-5296, 610-869-4024, FAX:610-869-0171
cuserv@jacksonimmuno.com, www.jacksonimmuno.com
www.jireurope.com

Product Specifications

Product: Rabbit Anti-Alpaca^{††} Whole Serum

Code Number: 328-001-001

Lot Number: 148116

Physical State: Freeze-dried solid

Size: 2.0 ml

Protein Concentration: 35.8 mg/ml

Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6

Preservative: 0.05% Sodium Azide

Storage and Rehydration: Store freeze-dried solid at 2-8°C. Rehydrate with 2.0 ml dH₂O and centrifuge if not clear. Prepare working dilution on day of use. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid.

Extended Storage after Rehydration: Aliquot and freeze undiluted product at -20°C or below. Avoid repeated freezing and thawing.
Expiration date: one year from date of rehydration. The expiration date may be extended if test results are acceptable for the intended use.

Purity: The antibody was delipidated and ammonium sulfate fractionated from whole antiserum.

Antibody Specificity: Antisera are obtained by injecting host animals with whole serum.

Country of Origin: USA

Note: For *in vitro* research use only, not for diagnostic or therapeutic use. This product is not a medical device.

††Warning: Bovine serum albumin (BSA) and dry milk may contain IgG. Secondary antibodies directed against alpaca, bovine, goat, horse, or sheep react with bovine IgG. Therefore, use of BSA or milk to block or dilute these antibodies may significantly increase background and/or reduce antibody titer. For blocking, normal serum (5% v/v) from the host species of the labeled secondary antibody is recommended.